

11-14-2014

Prescription Drug Abuse and Misuse in Southern Appalachia: An Epidemiologic Perspective

Billy Brooks

East Tennessee State University, brooksb1@etsu.edu

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>



Part of the [Substance Abuse and Addiction Commons](#)

Citation Information

Brooks, Billy. 2014. Prescription Drug Abuse and Misuse in Southern Appalachia: An Epidemiologic Perspective. *Controversies in Pain Management. Opioids: On the Horns of a Dilemma*, Johnson City, TN. <https://www.etsu.edu/com/cme/controversiesinpain.php>

This Presentation is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.


Prescription Drug Abuse and Misuse in Southern Appalachia: An Epidemiologic Perspective

Prescription Drug Abuse and Misuse in Southern Appalachia: An Epidemiologic Perspective

Billy Brooks, DrPH(c), MPH
Department of Biostatistics and
Epidemiology



Drug overdose death rates in the US have **more than tripled** since 1990.⁵



*Deaths are those for which poisoning by drugs (illicit, prescription, and over-the-counter) was the underlying cause.

National Vital Statistics System. Drug overdose death rates by state, 2008.

In 2008, there were 14,800 prescription painkiller deaths.⁴

For every **1** death there are...



10 treatment admissions for abuse⁹

32 emergency dept visits for misuse or abuse⁶

130 people who abuse or are dependent⁷


825
nonmedical
users⁷

12,210,000


CDC. Vital Signs: Overdoses of Prescription Opioid Pain Relievers—United States, 1999–2008. MMWR 2011; 60: 1–6

Substance Abuse and Mental Health Services Administration. Results from the 2010 National Survey on Drug Use and Health: volume 1: summary of national findings. Rockville, MD: Substance Abuse and Mental Health Services Administration, Office of Applied Studies; 2011. Available from URL: <http://oas.samhsa.gov/2k10/NSDUH/2k10NSDUH/2k10Results.htm#2.16>

Past Year Initiates of Illicit Drugs, 12 or Older, 2012



Drug overdose deaths of all intents by major drug type, U.S., 1999-2009



Source: National Vital Statistics System. The reported 2009 numbers are underestimates. Some overdose deaths were not included in the total for 2009 because of delayed reporting of the final cause of death.

Prescription vs. Illicit (2011)

- Drug overdose leading cause of injury death
- 80% (33,071) of overdose deaths unintentional
- 58.3% (1.4 million) of overdose ED Admissions related to pharmaceuticals
- 55% (22,810) of drug overdose deaths related to pharmaceuticals



Non-medical Prescription Drug Use (NMPDU)

“...the use of a medication without a prescription, in a way other than as prescribed, or for the experience or feelings elicited...”


-National Institute on Drug Abuse




COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Drug Overdose Rates by State, 2008




Percent Change in Unintentional Poisoning Mortality Rates, by Rural Status of State – United States, 1999-2004



Centers for Disease Control and Prevention. (2007). Unintentional poisoning deaths -- united state, 1999-2004. Morbidity and Mortality Weekly Report, 56(5), 93-96.

Drug Overdose Death Rates Are Increasing

**Death Rates from Drug Overdoses
per 100,000 population, Tennessee vs. US**



Source: Office of Policy, Planning and Assessment, Tennessee Department of Health – Death Certificates

Source: NCHS Data Brief, No. 81, December 2011, "Drug Poisoning Deaths in the United States, 1980 – 2008", Data table for Figure 1


Drug Dependence Surveillance System (Week 52)¹

Reporting Summary (Week 52)


Source of Maternal Exposure	Maternal County of Residence (By Health Department Region)	# Cases	% Cases	Number of cases reported		
				# Cases ²	% Cases	
Supervised replacement	Davidson	34	4.0%	397	46.4%	
Supervised pain treatment	East	240	28.1%	165	19.3%	
Therapy for psychological	Hamilton	17	2.0%	67	7.8%	
Prescription substances	Jackson/Madison	2	0.2%	341	39.9%	
Non-prescription substances	Knox	99	11.6%	236	27.6%	
No known exposure	Mid-Cumberland	57	6.7%	11	1.3%	
No response	North East	124	14.5%	19	2.2%	
	Shelby	18	2.1%			
	South Central	27	3.2%			
	South East	12	1.4%			
	Sullivan	84	9.8%			
	Upper Cumberland	110	12.9%			
	West	31	3.6%			
	Total	855	100%			

1. Summary reports are archived.
2. Multiple maternal substance use reports are included.


**Table I-1. Drug Dependent Newborns (Neonatal Abstinence Syndrome)
Surveillance Summary 2013**




Deaths by Accidental Poisonings, 2002-2011, Region 1 vs. TN



Drug Poisoning Mortality Rate, by Region 1 Counties, 2004-2010



What's Driving NMPDU?



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Supply



COLLEGE *of*
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY



46 



Each day, 46 people die from an overdose of prescription painkillers* in the US.

 **259 M**


Health care providers wrote 259 million prescriptions for painkillers in 2012, enough for every American adult to have a bottle of pills.

10 

10 of highest prescribing states for painkillers are in the South.




Total Number of Opioid Prescriptions Dispensed by U.S. Retail Pharmacies, 1991–2010




Source: SDI's Vector One®: National (VONA)


Opioid Prescription Rates by County—TN, 2007




Prescription Rate per 100 Population




Opioid Prescription Rates by County—TN, 2008




Prescription Rate per 100 Population




Opioid Prescription Rates by County—TN, 2009




Prescription Rate per 100 Population




Opioid Prescription Rates by County—TN, 2010




Prescription Rate per 100 Population



Opioid Prescription Rates by County—TN, 2011



Prescription Rate per 100 Population



Drug Dispensing in Tennessee



51 pills
per every
Tennessean
over age 12

275.5 Million Hydrocodone Pills



22 pills
per every
Tennessean
over age 12

116.6 Million Xanax Pills




21 pills
per every
Tennessean
over age 12

113.5 Million Oxycodone Pills



Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older: 2011-2012



¹ The Other category includes the sources "Wrote Fake Prescription," "Stole from Doctor's Office/Clinic/Hospital/Pharmacy," and "Some Other Way."

TABLE 5
Source of Prescription Drugs^a among Those Who Used in Last Year
Grade 12, 2007–2011
 (Entries are percentages.)

Where did you get the [insert drug name here] you used without a doctor's orders during the past year? (Mark all that apply.)


	<u>Amphetamines</u>		<u>Tranquilizers</u>		<u>Narcotics other than Heroin</u>	
	<u>2007–2008</u>	<u>2009–2011</u>	<u>2007–2008</u>	<u>2009–2011</u>	<u>2007–2008</u>	<u>2009–2011</u>
Bought on Internet	4.6	4.5	2.4	5.2	2.3	1.6
Took from friend/relative without asking	19.6	14.2	21.1	22.9	24.2	21.8
Took from a friend	—	4.9	—	6.6	—	4.7
Took from a relative	—	9.3	—	16.3	—	17.2
Given for free by friend or relative	58.2	66.4	59.8	78.3	50.5	70.2
Given for free by a friend	—	5.5	—	5.9	—	5.9
Given for free by a relative	—	9.9	—	24.4	—	19.9
Bought from friend or relative	45.0	41.6	44.1	51.7	37.1	41.4
Bought from a friend	—	4.9	—	4.9	—	3.4
Bought from a relative	—	2.6	—	6.8	—	5.1
From a prescription I had	15.1	18.1	18.4	14.7	40.2	34.7
Bought from drug dealer/stranger	26.7	21.8	24.2	26.7	18.6	16.4
Other method	17.8	13.8	7.5	9.9	8.5	10.5
<i>Weighted N =</i>	261	394	226	289	361	447

Source. The Monitoring the Future study, the University of Michigan.


Note. '—' indicates data not available.

^aIn 2009, the response categories were expanded to differentiate between friends and relatives.

Diversion Tree




Rates of prescription painkiller sales, deaths and substance abuse treatment admissions (1999-2010)



SOURCES: National Vital Statistics System, 1999-2008; Automation of Reports and Consolidated Orders System (ARCOS) of the Drug Enforcement Administration (DEA), 1999-2010; Treatment Episode Data Set, 1999-2009

What's Driving NMPDU?



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Demand/ Potential Demand



COLLEGE *of*
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

In 2008, there were 14,800 prescription painkiller deaths.⁴

For every **1** death there are...



10 treatment admissions for abuse⁹

32 emergency dept visits for misuse or abuse⁶

130 people who abuse or are dependent⁷

825 nonmedical users⁷

CDC. Vital Signs: Overdoses of Prescription Opioid Pain Relievers—United States, 1999–2008. MMWR 2011; 60: 1–6

Substance Abuse and Mental Health Services Administration. Results from the 2010 National Survey on Drug Use and Health: volume 1: summary of national findings. Rockville, MD: Substance Abuse and Mental Health Services Administration, Office of Applied Studies; 2011. Available from URL: <http://oas.samhsa.gov/2k10NSDUH/2k10NSDUHResults.htm#2.16>



13.3%

3.3%

KEY

- Only require screening
- Require brief intervention
- Require referral to treatment

Source: SAMHSA funded
MASBIRT program, N=173,714



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Individual-Level Determinants


- Age
- Gender
- Social/ Emotional competence
- Mental health
- Educational Attainment
- Employment Status
- Marital status
- Housing
- History of trauma
- Age of drug use initiation



COLLEGE of
PUBLIC HEALTH


EAST TENNESSEE STATE UNIVERSITY

Past Year Nonmedical Use of Pain Relievers, by Detailed Age Category



SOURCE: <http://www.oas.samhsa.gov/nhsda/>

Prescription painkiller overdose deaths are a growing problem among women.



Socio-Familial Determinants

- Social-network size and characteristics
- Parental involvement
- Peer drug use
- Familial support/ bonding
- Divorced parents



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Adverse Childhood Experiences

	Number	95% CI			Adj.	95% CI	
	Resp.*	Wt. %	LL	UL	Odds Ratio [†]	LL	UL
Misuse of Prescription Pain Medications							
0 ACEs	3,815	1.4	0.9	2.1	1.00	Referent	
1 to 3 ACEs	3,679	4.1	3.2	5.1	2.76	1.62	4.70
4 or more ACEs	1,232	9.6	7.2	12.6	6.25	3.49	11.20



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Neighborhood-Level Determinants

- Unemployment rate
- Average educational attainment
- Drug arrest rate
- Average income
- Community cohesion
- Diversion control (e.g. drug drop boxes, PDMPs)



Opioid-Related Mortality

Prescriber Behavior


- High volume prescribing
- Sales
- Dosage
- Oxycodone prescribing
- Methadone prescribing



COLLEGE of
PUBLIC HEALTH


EAST TENNESSEE STATE UNIVERSITY

Growth of methadone use for pain and methadone overdoses



Sources: National Vital Statistics System, 1999-2009; Drug Enforcement Administration Automation of Reports and Consolidated Orders System (ARCOS), 1999-2010.

Death rate from overdoses caused by a single prescription painkiller



Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Statistics and Quality, Drug Abuse Warning Network Medical Examiner Component, 2009.

Opioid-Related Mortality

Individual Determinants


- SA history
- Diversion
- Doctor shopping
- Drug substitution
- Polydrug toxicity
- Socio-demographics



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Number of High Utilization Patients by Quarter in TN CSMD, 2010-2013



Opioid-Related Mortality


Environmental Determinants

- Urbanization
- Geography
- Public policy
- Intervention
- Media
- PDMP



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY




What Can be Done?



COLLEGE *of*
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY


What's Driving NMPDU?



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Aggregate Efforts



Prescribing Guidelines

Five studies in different states have shown that ~10-15% of prescribers prescribe ~65-80% of OPRs



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Prescribing Guidelines

MED: Morphine Equivalent Dose;
method of standardizing the volume of
consumed opioids in a day

Strong epidemiologic evidence for a
significant increase in opioid related morbidity
and mortality above 100-120 mg/d MED

MED. Morphine Equivalent

Dosage

- Study of 45 overdoses
- Hazard ratio:
 - 50-100 mg/d MED = almost 4x increase in risk
 - >100 mg/d MED = almost 9x increase in risk

Dunn et al. 2010. Ann Int Med 152; 85-92

Table 3. Hazard Ratios Between Recent Opioid Doses and Overdose*


Opioid Dose	Patients Who Overdosed, n	Person-Years	Overdose Rate (95% CI) per 100 000 Person-Years	Hazard Ratio for All Overdose Events (95% CI)†	Hazard Ratio for Serious Overdose Events (95% CI)†‡
None	6	16 780	36 (13–70)	0.31 (0.12–0.80)	0.19 (0.05–0.68)
1 to <20 mg/d				1.00	1.00
20 to <50 mg/d	6	2311	260 (95–505)	1.44 (0.57–3.62)	1.19 (0.40–3.60)
50 to <100 mg/d				3.73 (1.47–9.50)	3.11 (1.01–9.51)
≥100 mg/d				8.87 (3.99–19.72)	11.18 (4.80–26.03)
Any opioid use	45	17 582	256 (187–336)	5.16 (2.14–12.48)	8.39 (2.52–27.98)

* Opioid-related overdose death or nonfatal event.

† Adjusted for smoking, depression, substance abuse, comorbid conditions, pain site, age, sex, recent sedative-hypnotic prescription, and recent initiation of opioid use.

‡ Opioid-related overdose death or serious nonfatal event (n = 40).

Opioid prescribing guidelines



Higher dose opioids= >/+ 120 mg/day MED


Controlling the Supply




COLLEGE *of*
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Status of State Prescription Drug Monitoring Programs (PDMPs)






Painkillers And The Heroin Market

A growing number of people are using heroin in recent years, in part because it can be cheaper and easier to find than opioid painkillers purchased on the black market. Most heroin users were first hooked on prescription opioids, which generated \$11 billion in 2010 for the pharmaceutical industry.

Substance abuse treatment facilities admissions by primary drug



*2012 data for Mississippi, Pennsylvania, and West Virginia are not available.

4 out of 5

new heroin users have abused painkillers.




A Cheaper High

\$30 can buy one oxycodone pill on the street in New York...




or six hits of heroin.




PDMP Training & Technical Assistance Center

Prescription Drug Monitoring Programs (PDMPs) Interstate Data Sharing Status



National Take Back Day: **October 26, 2013**


Total Weight Collected (pounds): 647,211 (324 Tons)



Drug Enforcement Administration
Diversion Control Program

Permanent Collection Boxes

Figure II-2. Number of Permanent Prescription Drug Collection Boxes




Source: Tennessee Department of Mental Health and Substance Abuse Services (2013)

Breaking News

D.E.A. to Allow Return of Unused Pills to Pharmacies

By CATHERINE SAINT LOUIS SEPT. 8, 2014

 Share

 Tweet

 Pin

 Email

 Save

MORE >



Concerned by rising rates of prescription drug abuse, the Drug Enforcement Administration announced Monday that it would permit consumers to return unused prescription medications like opioid painkillers to pharmacies.

The move is intended to help reduce stockpiles of unneeded medicines in homes, which are often pilfered by teenagers. Under the new regulation, patients and their relatives will also be allowed to mail unused prescription drugs to an authorized collector using packages to be made available at pharmacies and other locations, like libraries and senior centers.

The new regulation, which will go into effect in a



Don't miss a single story. Try a Times digital subscription for 99¢

ALREADY A SUBSCRIBER? LOG IN.




COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

The Ambulance or the Fence?

Prevention Approach
Usual Approach



Primary Prevention Works

- Parenting programs work:
 - Parental monitoring
 - Authoritative parenting
- Parents need to be engaged
- Kids at risk need to be trained
- We can scale up effective programs to prevent uptake of NMPDU



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

NREPP is a searchable online registry of [more than 300 interventions](#) supporting mental health promotion, substance abuse prevention, and mental health and substance abuse treatment. We connect members of the public to intervention developers so they can learn how to implement these approaches in their communities.

NREPP is not an exhaustive list of interventions, and inclusion in the registry does not constitute an endorsement. [Learn More >](#)

Basic Search

Advanced Search

View All

Interventions

Find an Intervention

Search

Find interventions reviewed by NREPP.

News

 [Learn About NREPP's RSS Feed](#)

New Intervention Summary Available - 09/23/2013

Read the newly posted summary for *Interactive Journaling*

[Read more >](#)

New Intervention Summary Available - 09/23/2013

Read the newly posted summary for *Family Wellness: Survival Skills for Healthy Families*

[Read more >](#)



Subscribe

Enter your email address to receive monthly NREPP updates.

300th Intervention Summary Posted

SAMHSA's NREPP reached a new milestone, publishing its 300th summary of an evidence-based substance abuse or mental health intervention. See the [SAMHSA Bulletin](#) for more information about NREPP and this milestone.

Early Intervention

Screening, Brief Intervention, and Referral to Treatment (SBIRT)



Naloxone




- Reverses overdose by blocking receptors
- White House (ONDCP) and Attorney General encourage first responders to carry
- In conjunction with “Good Samaritan” law can effectively reduce mortality rate




COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Number of Deaths per 100,000 Population from Unintentional Drug Poisoning in Wilkes County, NC, 2004-2011



States with Naloxone Access and Overdose Good Samaritan Laws as of April 2014



Source: The Network for Public Health Law, *Legal Interventions to Reduce Overdose Mortality: Naloxone Access and Overdose Good Samaritan Laws*, web document, https://www.networkforphl.org/_asset/qz5pvn/network-naloxone-10-6.pdf, accessed March 15, 2014 and the Georgia governor's website.

*Law in Alaska explicitly requires courts to take the fact that a good samaritan summoned medical assistance into account at sentencing.

Northeast TN Resources

Generation Rx

<http://etsugenerationrxprovidertoolkit.weebly.com/prescriber-resources.html>

Prescription Drug Abuse and Misuse Working Group

<http://www.etsu.edu/cph/pdam/>



COLLEGE of
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

Questions?



COLLEGE *of*
PUBLIC HEALTH

EAST TENNESSEE STATE UNIVERSITY

References

1. Centers for Disease Control and Prevention. (2014). Prescription drug overdose in the united states: Fact sheet. Retrieved from <http://www.cdc.gov/homeandrecreationalafety/overdose/facts.html>.
2. TN Department of Mental Health and Substance Abuse Services. Prescription for Success. Retrieved from <http://tn.gov/mental/prescriptionforsuccess/Prescription%20For%20Success%20SECTION%201.pdf>. Sept. 2014.
3. United States Department of Health and Human Services (USDHHS), Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Compressed Mortality File (CMF) on CDC WONDER Online Database.
4. King, N. B., Fraser, V., Boikos, C., Richardson, R., & Harper, S. (2014). Determinants of increased opioid-related mortality in the united states and canada, 1990-2013: A systematic review. *American Journal of Public Health*, 104(8), e32-42. doi:10.2105/AJPH.2014.301966 [doi]
5. Swendsen, J., Conway, K. P., Degenhardt, L., Glantz, M., Jin, R., Merikangas, K. R., . . . Kessler, R. C. (2010). Mental disorders as risk factors for substance use, abuse and dependence: Results from the 10-year follow-up of the national comorbidity survey. *Addiction (Abingdon, England)*, 105(6), 1117-1128. doi:10.1111/j.1360-0443.2010.02902.x [doi]
6. Brason F. Project Lazarus: An Innovative Community Response to Prescription Drug Overdose. *NC Med J.* 2013; 74(3):259-261.
7. Wang, K. H., Becker, W. C., & Fiellin, D. A. (2013). Prevalence and correlates for nonmedical use of prescription opioids among urban and rural residents. *Drug and Alcohol Dependence*, 127(1-3), 156-162. doi:10.1016/j.drugalcdep.2012.06.027 [doi]

